

Noncommutativity in quantum cosmology and the Hierarchy problem

F. Darabi^{*}, A. Rezaei-Aghdam[†], A. R. Rastkar

Department of Physics, Azarbaijan University of Tarbiat Moallem, 53714-161, Tabriz, Iran .

February 7, 2008

Abstract

We study the quantum cosmology of an empty (4+1)-dimensional Kaluza-Klein cosmology with a negative cosmological constant and a FRW type metric with two scale factors, one for 4-D universe and the other for one compact extra dimension. By assuming the noncommutativity in the corresponding mini-superspace we suggest a solution for the Hierarchy problem, at the level of Wheeler-DeWitt equation.

PACS: 98.80.Qc

^{*}e-mail: f.darabi@azaruniv.edu

[†]e-mail: rezaei-a@azaruniv.edu

Recently, the study of various physical theories from noncommutative point of view, such as string theory [1], quantum field theory (see for instance [2]), quantum mechanics [3], and classical mechanics [4], has been of particular interest. In particular, a new interest has been developed to study the noncommutative *quantum cosmology* [5], [6], [7]. In these studies, the influence of noncommutativity at early universe was explored by the formulation of a version of noncommutative quantum cosmology in which a deformation of *mini-superspace* is required instead of space-time deformation [5].

We know that in the study of homogeneous universes, the metric depends only on the time parameter. Thus, one can find a model with a finite dimensional configuration space, the so-called mini-superspace, whose variables are the three-metric components. The quantization of these models can be performed by using of the rules of quantum mechanics. The mini-superspace construction is a procedure to define quantum cosmological models in the search for describing the quantum features of the early universe. The influence of noncommutativity on the mini-superspace has already been considered for the cosmological models with Kantowski-Sachs and FRW metrics [5], [6]. Here, we consider the effect of noncommutativity on the configuration space in the model which was previously studied by one of the authors [8]. This model shows an empty (4+1) dimensional Kaluza-Klein universe with a negative cosmological constant and a FRW type metric having two scale factors. Following the idea in regarding the Wheeler-Dewitt equation more primitive than the classical Einstein equations ¹ we study the noncommutativity in the quantum cosmology of this model and try to propose a solution for the Hierarchy problem, in the context of quantum cosmology.

The experimental upper bound on the value of cosmological constant is extremely small. On the other hand, it is usually assumed that an *effective* cosmological constant describes the energy density of the vacuum $\langle \rho_{vac} \rangle$. In fact, the vacuum energy density $\langle \rho_{vac} \rangle$ is a quantum field theory contribution to the effective cosmological constant

$$\Lambda_{eff} = \lambda + \kappa \langle \rho_{vac} \rangle,$$

where λ is a small bare cosmological constant. The calculations show that, these contributions affect enormously the value of effective cosmological constant as $\langle \rho_{vac} \rangle \sim M_P^4$, where M_P is the Planck mass which defines the ultraviolet cutoff scale of the quantum gravity. This is the well-known cosmological constant problem. However, there is another fundamental energy scale in nature, namely the electroweak scale M_{EW} whose experimental investigation is of particular interest and the corresponding interaction has been probed successfully. Over the past two decades there was a great interest to explain the smallness of M_{EW}/M_P , which is known as the Hierarchy problem. Recently, a great amount of interest has been concentrated on solving this problem based on the existence of large extra dimensions [9]. Some attempts have also been done to solve this problem based on the noncommutativity in the space-time coordinates [10]. To the authors knowledge, this problem has not yet been paid attention by considering the noncommutativity in the mini-superspace coordinates of a quantum cosmology.

In this paper, we introduce a mechanism based on the existence of noncommutativity in the mini-superspace of a quantum cosmology corresponding to an empty (4 + 1)-dimensional Kaluza-Klein cosmology. It is to be noted that the present mechanism does not claim to solve

¹This idea was followed by Hawking and Page investigating the wormholes [11]

the problem in a fundamental way because we just solve the problem in a special and simple model. However, this may shed light on the similar approaches to solve the problem in a more fundamental way.

We start with the metric considered in [8] in which the space-time is assumed to be of FRW type which has a compact space, namely the circle S^1 . We adopt the chart $\{t, r^i, \rho\}$ with t , r^i and ρ denoting the cosmic time, the space coordinates and the compact space coordinate, respectively. We therefore take

$$ds^2 = -dt^2 + R^2(t) \frac{dr^i dr^i}{(1 + \frac{kr^2}{4})^2} + a^2(t) d\rho^2, \quad (1)$$

where $k = 0, \pm 1$ and $R(t)$, $a(t)$ are the scale factors of the universe and compact dimension, respectively. The curvature scalar corresponding to metric (1) is obtained as

$$\mathcal{R} = 6 \left[\frac{\ddot{R}}{R} + \frac{k + \dot{R}^2}{R^2} \right] + 2 \frac{\ddot{a}}{a} + 6 \frac{\dot{R}}{R} \frac{\dot{a}}{a}, \quad (2)$$

where a dot represents differentiation with respect to t . Substituting this result into Einstein-Hilbert action with a cosmological constant Λ

$$I = \int \sqrt{-g} (\mathcal{R} - \Lambda) dt d^3r d\rho, \quad (3)$$

and integrating over spatial dimensions gives an effective Lagrangian L in the mini-superspace (R, a) as

$$L = \frac{1}{2} R a \dot{R}^2 + \frac{1}{2} R^2 \dot{R} \dot{a} - \frac{1}{2} k R a + \frac{1}{6} \Lambda R^3 a. \quad (4)$$

By defining $\omega^2 \equiv -\frac{2\Lambda}{3}$ and changing the variables as

$$u = \frac{1}{\sqrt{8}} \left[R^2 + R a - \frac{3k}{\Lambda} \right], \quad v = \frac{1}{\sqrt{8}} \left[R^2 - R a - \frac{3k}{\Lambda} \right], \quad (5)$$

L takes on the form

$$L = \frac{1}{2} \left[(\dot{u}^2 - \omega^2 u^2) - (\dot{v}^2 - \omega^2 v^2) \right]. \quad (6)$$

The assumption that the full (4+1) dimensional Einstein equations hold, implies that the Hamiltonian corresponding to L in (6) must vanish, that is

$$H = \frac{1}{2} \left[(\dot{u}^2 + \omega^2 u^2) - (\dot{v}^2 + \omega^2 v^2) \right] = 0, \quad (7)$$

which describes an isotropic oscillator-ghost-oscillator system.

The corresponding quantum cosmology is described by the Wheeler-DeWitt equation resulting from Hamiltonian (7) and can be written as

$$\{[p_u^2 + \omega^2 u^2] - [p_v^2 + \omega^2 v^2]\} \Psi(u, v) = 0. \quad (8)$$

Now, we consider the effect of noncommutativity on the configuration space in the above model. The noncommutative quantum mechanics is defined by the following commutators [3]

$$[u, v] = i\theta, \quad [u, p_u] = i, \quad [v, p_v] = i, \quad [p_u, p_v] = 0, \quad (9)$$

where we use the natural units ($\hbar = 1$). The corresponding noncommutative Wheeler-DeWitt equation can be written by use of the star product as [3], [5]

$$H * \Psi = 0. \quad (10)$$

We can always represent the noncommutative harmonic oscillator in terms of anisotropic harmonic oscillator in the commuting coordinates [3]

$$[u, v] = 0, \quad [u, p_u] = i, \quad [v, p_v] = i, \quad [p_u, p_v] = 0. \quad (11)$$

Therefore, we obtain anisotropic oscillator-ghost-oscillator Wheeler-DeWitt equation

$$\left\{ \left[\frac{p_u^2}{4} + \omega^2 \left(u - \frac{\theta p_v}{2} \right)^2 \right] - \left[\frac{p_v^2}{4} + \omega^2 \left(v + \frac{\theta p_u}{2} \right)^2 \right] \right\} \Psi(u, v) = 0, \quad (12)$$

where the transformations $u \rightarrow u - \theta p_v$ and $v \rightarrow v + \theta p_u$ have been used. However, in two dimensions we can also represent the two-dimensional anisotropic oscillator-ghost-oscillator as the two-dimensional isotropic oscillator-ghost-oscillator in the presence of an effective magnetic field as follows

$$\{[(p_u - A_u)^2 + \omega'^2 u^2] - [(p_v - A_v)^2 + \omega'^2 v^2]\} \Psi(u, v) = 0, \quad (13)$$

where

$$\begin{cases} \omega'^2 \equiv \frac{4\omega^2}{(1-\omega^2\theta^2)^2}, \\ A_u \equiv \frac{2\omega^2\theta}{1-\omega^2\theta^2}v, \\ A_v \equiv -\frac{2\omega^2\theta}{1-\omega^2\theta^2}u. \end{cases} \quad (14)$$

The above equation can be written in a more convenient form

$$\left\{ \left[p_u^2 + \left(\omega'^2 - \frac{B^2}{4} \right) u^2 \right] - \left[p_v^2 + \left(\omega'^2 - \frac{B^2}{4} \right) v^2 \right] + B(vp_u + up_v) \right\} \Psi(u, v) = 0, \quad (15)$$

where $B = -\frac{4\omega^2\theta}{1-\omega^2\theta^2}$ is derived through $\mathbf{B} = \nabla \times \mathbf{A}$ from

$$A_u = -\frac{B}{2}v, \quad A_v = \frac{B}{2}u. \quad (16)$$

The B -term in Eq.(15) deserves more scrutiny so that we can find the correct frequency for this system. To this end, it is useful to compare this oscillator-ghost-oscillator system (15)

with the corresponding oscillator-oscillator system. By straightforward calculations we find for the latter system

$$H = \frac{1}{2} \sum_{i=1,2} [(p_i - A_i)^2 + \omega^2 x_i^2] = \frac{1}{2} \sum_{i=1,2} [p_i^2 + (\omega^2 + \frac{B^2}{4}) x_i^2] - \frac{1}{2} B(x_1 p_2 - x_2 p_1), \quad (17)$$

where $A = (-x_2, x_1, 0) \frac{B}{2}$. The B -term in Eq.(17) is the magnetic potential energy and $\sqrt{(\omega^2 + \frac{B^2}{4})}$ is the effective frequency of the system. It is obvious that the B -term has an independent role and does not contribute to the oscillator frequency, at all. Now, comparing the two systems we realize that each term in Eq.(17) has a corresponding term in Eq.(15). For example, the first two brackets in Eq.(17) correspond to the first two brackets in Eq.(15), and the third B -term in Eq.(17) corresponds to the third B -term in Eq.(15). Therefore, the corresponding terms have the same roles with the difference that the terms in Eq.(15) have the *ghost* characters. This occurs in other examples, as well. For instance, the two-dimensional oscillator with no external magnetic field is defined by

$$H = (p_x^2 + p_y^2) + \omega^2(x^2 + y^2), \quad (18)$$

whereas the corresponding oscillator-ghost-oscillator system is defined by

$$H = (p_x^2 - p_y^2) + \omega^2(x^2 - y^2). \quad (19)$$

Here, the first parenthesis in both systems has the kinetic term role and the second parenthesis plays the role of potential term, with the difference that the terms in Eq.(19) have the ghost characters.

Therefore, as the B -term in Eq.(17) is an independent potential term which does not contribute to the effective frequency $\sqrt{(\omega^2 + \frac{B^2}{4})}$, the corresponding B -term in Eq.(15) should also be an independent potential term which does not contribute to the effective frequency, as well.

In this regard, we can write

$$\tilde{\omega}^2 = \omega'^2 - \frac{B^2}{4} = \frac{4\omega^2}{(1 - \omega^2\theta^2)}, \quad \omega^2\theta^2 \leq 1, \quad (20)$$

which defines the resultant frequency of the isotropic oscillator-ghost-oscillator in the presence of a constant magnetic field B . This frequency comes from a ghost-like combination of the oscillator term ω'^2 and the cyclotron term $\frac{B^2}{4}$.

In Eq.(8), the oscillator frequency ω was defined by the effective cosmological constant $\Lambda_{eff} = -\frac{3}{2}\omega^2$. Comparing with Eq.(15), we suppose $\tilde{\omega}$ to be defined by a new effective cosmological constant $\tilde{\Lambda}_{eff} = -\frac{3}{2}\tilde{\omega}^2$. Substituting these definitions into Eq.(20) we obtain

$$\tilde{\Lambda}_{eff} = \frac{4\Lambda_{eff}}{(1 - \frac{2}{3}\theta^2|\Lambda_{eff}|)}. \quad (21)$$

This is a redefinition of the effective cosmological constant due to the noncommutativity. However, we know that the effective cosmological constant, in principle, is a measure of the

ultraviolet cutoff in the theory. Therefore, the above equation can also be considered as a redefinition of the cutoff in the theory due to the noncommutativity. This is the main result which is considered to solve the Hierarchy problem in the present model. To this end, we take

$$\theta^2 = \frac{3 M_P^4 - 4 M_{EW}^4}{2 M_P^4 M_{EW}^4}, \quad (22)$$

where M_{EW} is the electroweak mass scale. Moreover, we take $\Lambda_{eff} \sim M_{EW}^4$ representing M_{EW} as the natural cutoff in the original commutative model. Therefore, we obtain from the above equation

$$\tilde{\Lambda}_{eff} \sim M_P^4, \quad (23)$$

which defines the cutoff in the noncommutative model. In other words, if we assume M_{EW} to be the natural cutoff in the original commutative model, the Planck mass is then the cutoff in the noncommutative model. This solves the Hierarchy problem at the level of quantum Wheeler-DeWitt equation by assuming that M_{EW} is the only fundamental mass scale in the model, and M_P is the mass scale which is appeared due to introducing the noncommutativity in the mini-superspace. Put another way, one may suppose that the universe, in principle, has just one fundamental energy scale for all interactions, namely M_{EW} . The quantum gravity sector of this universe must then be described by the Wheeler-DeWitt equation (8) with the vacuum energy density of the same scale $\omega^2 = -\frac{2}{3}\Lambda_{eff} \sim M_{EW}^4$. However, the energy scale of the quantum gravity which we expect to experience in the universe is defined by the Planck mass M_P and not M_{EW} . This discrepancy between M_P and M_{EW} , namely the Hierarchy problem is solved by the assumption of noncommutativity in the quantum gravity sector of the universe which leads Eq.(8) to Eq.(15) with the vacuum energy density of the Planck scale $\tilde{\omega}^2 = -\frac{2}{3}\tilde{\Lambda}_{eff} \sim M_P^4$. It then turns out that M_P is not a new fundamental scale and its enormity $M_P \gg M_{EW}$ is simply a consequence of the noncommutativity in the quantum gravity sector of the model, namely the Wheeler-DeWitt equation.

It is worth noting that having the exact numeric coefficients in Eq.(22) is very important to get Eq.(23) which solves the Hierarchy problem. This in particular means that our proposed solution for the Hierarchy problem comes with a *fine tuning*. Although this fine tuning makes the proposed solution less appealing, however, it is still a novel method, as the Hierarchy problem seems to be solved with tuning only one parameter, θ .

Acknowledgment

We would like to thank the referee very much for the useful and constructive comments. This work has been financially supported by the Research Department of Azarbaijan University of Tarbiat Moallem, Tabriz, Iran.

References

- [1] N. Seiberg and E. Witten, JHEP **09** (1999), 032.
- [2] R. Szazbo, Phys. Rep **378** (2003), 207.

- [3] M. Chaichian, M. M. Sheikh-Jabbari, and A. Tureanu, Eur. Phys. J. C **36** (2004) 251; L. Mezincescu, *Star operation in quantum mechanics*, hep-th/0007046; J. Gamboa, M. Loewe, and J. c. Rojas, Phys. Rev. D **64** (2001), 067901; S. Bellucci and A. Nersessian, Phys. Lett. B **542** (2002) 295.
- [4] J. M. Romero, J. A. Satiago, and D. Vergara, Phys. Lett. A **310** (2003), 9; A. E. F. Djemai, *On noncommutative classical mechanics*, hep-th/0309034.
- [5] H. García-Compeán, O. Obregón, and C. Ramírez, Phys. Rev. Lett **88** (2002), 161301.
- [6] G. D. Barbosa and N. Pinto-Neto, Phys. Rev. D **70** (2004), 103512.
- [7] G. D. Barbosa, Phys. Rev. D **71** (2005), 063511.
- [8] F. Darabi and H. R. Sepangi, Class. Quantum. Grav **16** (1999), 1656.
- [9] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Lett. B **429** (1998), 263; L. Randall, R. Sundrum, Phys. Rev. Lett. **83** (1999), 3370.
- [10] F. Lizzi, G. Mangano, and G. Miele, Mod. Phys. Lett. A **16** (2001), 1; Xiao-Jun Wang, *Cosmological Constant as Vacuum Energy Density of Quantum Field Theories on Non-commutative Spacetime*, hep-th/0411248.
- [11] S. W. Hawking, D. N. Page, Phys. Rev. D **42** (1990), 2655.